

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 88311798.8

(22) Date of filing: 14.12.88

(51) Int. Cl.4: C07D 209/42 , C07D 471/04 ,
C07D 215/56 , C07D 241/44 ,
C07D 207/34 , C07D 209/18 ,
C07D 209/08 , A61K 31/40 ,
A61K 31/47 , A61K 31/495 ,
A61K 31/44 , //(C07D471/04,
221:00,209:00),(C07D471/04,
235:00,221:00)

(30) Priority: 15.12.87 US 132373

(43) Date of publication of application:
21.06.89 Bulletin 89/25

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE


(71) Applicant: PFIZER INC.
235 East 42nd Street
New York, N.Y. 10017(US)

(72) Inventor: Rosati, Robert Louis
RD 3, Box 66A Deans Mill Road
Stonington Connecticut(US)

(74) Representative: Moore, James William, Dr.
Pfizer Limited Ramsgate Road
Sandwich Kent CT13 9NJ(GB)

(54) Non-peptidic renin inhibitors.

(57) Simple amides and derivatives thereof useful for inhibiting the angiotensinogen-cleaving action of the enzyme renin of the formula:



wherein R₁ is (C₆-C₈)cycloalkyl or i-propyl; R₂ is (C₃-C₅)alkyl, phenyl, methylvinyl, dimethylvinyl, halovinyl, hydroxy(C₁-C₃)alkyl or amino(C₁-C₄)alkyl; R₃ is (C₁-C₆)alkyl or morpholinoethyl; and Het is defined to include certain mono and bicyclic nitrogen-containing heterocyclic rings or heterocyclalkyl groups.

NON-PEPTIDIC RENIN INHIBITORS

This invention relates to novel simple amides and derivatives thereof useful as antihypertensive agents.

The proteolytic enzyme renin, which has a molecular weight of about 40,000, is produced in and secreted into the blood by the kidney. It is known to be active *in vivo* in cleaving the naturally-occurring plasma glycoprotein angiotensinogen, in the case of human angiotensinogen at the bond between the leucine (10th) and valine (11th) amino acid residues at the N-terminal end of the angiotensinogen:

Asp-Arg-Val-Tyr-Ile-His-Pro-Phe-His-Leu-Val-

1 2 3 4 5 6 7 8 9 10 11

Ile-His-Ser-Glu-


12 13 14 15

The circulating N-terminal decapeptide (angiotensin I) formed by the above cleaving action of renin is subsequently broken down by the body to an octapeptide known as angiotensin II. Angiotensin II is known to be a potent pressor substance, i.e., a substance that is capable of inducing a significant increase in blood pressure and is believed to act by causing the constriction of blood vessels and the release of the sodium-retaining hormone aldosterone from the adrenal gland. Thus, the renin-angiotensinogen system has been implicated as a causative factor in certain forms of hypertension and congestive heart failure.

One means of alleviating the adverse effects of the functioning of the renin-angiotensinogen system is the administration of a substance capable of inhibiting the angiotensinogen-cleaving action of renin. A number of such substances are known including antirenin antibodies, pepstatin and naturally-occurring phospholipid compounds. European Patent Application No. 45,665 (published February 2, 1982) discloses a series of renin-inhibiting polypeptide derivatives of the formula

X-Y-Pro-Phe-His-A-B-Z-W

in which X may be hydrogen or an amino-protecting group, Y may be absent, B is a lipophilic amino acid residue, Z is an aromatic amino acid residue, W may be hydroxyl and A may be, inter alia,




with each of R¹ and R² being a lipophilic or aromatic side chain. According to the definitions set forth in this published patent application, it is not contemplated that either A or Z could be statine or that B could be lysine.


European Patent Application No. 77,028A (published April 20, 1983) discloses a series of renin-inhibiting polypeptide compounds having a non-terminal statine or statine derivative residue. Included within this series are compounds having a phenylalanine-histidine-statine sequence.

European Patent Application 132,304A also discloses the use of statine containing polypeptides as renin-inhibiting antihypertensive agents, and European Patent Application 114,993A discloses polypeptides containing cyclostatine, useful as renin-inhibiting antihypertensive agents.


The simple amide renin inhibitors, useful as antihypertensive agents, are of the formula



or a pharmaceutically acceptable salt thereof, wherein HET is




of a group of the formula



where X is hydrogen, (C₁-C₃)alkyl, (C₁-C₃)alkoxy, fluoro, chloro, bromo or cyano; Y is hydrogen, (C₁-C₃)alkyl, (C₁-C₃)alkoxy, fluoro or chloro; R₄ is hydrogen or (C₁-C₃)alkyl; n is an integer of 0 to 2; R₁ is (C₆-C₈)cycloalkyl or i-propyl; R₂ is (C₃-C₅)alkyl, phenyl, methylvinyl, dimethylvinyl, halovinyl, hydroxy(C₁-C₃)alkyl or amino(C₁-C₄)alkyl; and R₃ is (C₁-C₆)alkyl or morpholinoethyl.


Preferred within this group of compounds are those where HET is



R₁ is cyclohexyl and R₃ is methyl. Especially preferred are the compounds where X is 5-chloro, Y is hydrogen, R₄ is hydrogen, n is 0 and R₃ is either i-propyl or -C(Cl) = CH₂, and where X and Y are


hydrogen, R_4 is methyl, n is 0 and R_2 is *i*-propyl.

A second group of preferred compounds are those where HET is



R_1 is cyclohexyl and R_3 is methyl. Especially preferred is the compound where X is 5-chloro, Y is hydrogen, R_4 is hydrogen, n is 0 and R_2 is $-\text{CH}(\text{OH})\text{CH}_3$.

A third group of preferred compounds are those of the formula



where R_1 is cyclohexyl and R_2 is methylvinyl. Especially preferred are the compounds where X is 5-chloro, Y is hydrogen, R_4 is hydrogen, n is 0, R_2 is $-\text{CH}=\text{CHCH}_3$ and R_3 is methyl and where X is 5-chloro, Y is hydrogen, R_4 is hydrogen, n is 0, R_2 is $-\text{CH}=\text{CHCH}_3$ and R_3 is morpholinoethyl.

The present invention also includes a method for treating hypertension in a mammal which comprised administering to said mammal an antihypertensive effective amount of the compounds of the present invention and a pharmaceutical composition comprised of the compounds of the present invention and a carrier.

As previously indicated, the present invention embraces pharmaceutically acceptable salts of the biologically active compounds. Such salts are those which are non-toxic at the dosages administered. Since compounds of the invention may contain basic groups, acid addition salts are possible. Pharmaceutically acceptable acid addition salts include e.g. the hydrochloride, hydrobromide, hydroiodide, sulfate, bisulfate, phosphate, acid phosphate, acetate, lactate, maleate, mesylate, fumarate, citrate, acid citrate, tartrate, bitartrate, succinate, gluconate and saccharate salts.

The compounds of this invention exhibit antihypertensive activity *in vivo* in mammals, including humans. At least a substantial portion of this activity results from their ability to inhibit the cleavage of angiotensinogen by renin. Although we do not wish to be limited by the following theory of mechanism, it is likely that the mechanism of the renin-inhibiting activity of the compounds of the invention is their selective binding (as compared to angiotensinogen) to renin. The compounds of the invention exhibit an enzyme-inhibiting activity that is selective for renin. Because of their low molecular weights they exhibit favorable solubility characteristics in aqueous media and better absorbability, thus making oral administration feasible; and they can be synthesized at a commercially realistic cost. The compounds of the present invention are also useful against congestive heart failure.

The compounds of the invention may be prepared by methods familiar to those skilled in the art. The heterocyclic acids are coupled with the requisite amino lactone using coupling methods well known in the art for making amides. In this particular series of compounds the use of carbodiimides and *N*-hydroxybenzotriazole to form activated esters of the heterocyclic acids is preferred, although any of the many groups which are known to make activated esters can be employed. A second preferred means of coupling the requisite acid with the appropriate amino lactone is the use of diethyl cyanophosphate. This reagent is also well known for its use in the synthesis of amides.

Following the coupling of the appropriate heterocyclic acid and amino lactone, the lactone is treated with an amine resulting in the formation of the desired hydroxy-amide, having the necessary and desired stereo-chemistry to provide the antihypertensive activity which these compounds possess.

The starting reagents for the synthesis of these compounds are described herein or can be prepared by literature procedures. EPO application 86,305,995.2 [Publication 0212903(A2)] describes the preparation of polypeptides using an amino lactone similar to that described herein. Further, several of the heterocyclic

acids are available commercially.

The ability of these compounds to act as antihypertensive agents by the inhibition of renin can be shown by the following in vitro assay:

Inhibition of the Angiotensinogen-Cleaving Activity of Renin In Vitro

Blood plasma was obtained from healthy laboratory personnel, pooled and stored frozen until required. Before use, a quantity of this plasma was defrosted and centrifuged, and the supernatant mixed with protease inhibitors and buffered to pH 7.4. Renin inhibitors were added at different levels to different aliquots of the plasma supernatant, and the resulting mixtures (310 lambda) incubated for three hours at 37° C. along with renin inhibitor-free control mixtures. After incubation, the mixtures were quenched in ice water and each assayed for angiotensin I using angiotensin I antibody. The production of angiotensin I in the presence of a renin inhibitor was compared to that produced in the absence of the inhibitor, and a percentage inhibition was calculated. Using data obtained from duplicate incubations at each of several different inhibitor concentrations, the inhibitor concentration in the incubation mixture required to produce a fifty percent inhibition of the angiotensinogen-cleaving activity of renin, i.e., the IC₅₀ of the inhibitor, was calculated for various different inhibitors.

The angiotensin I in the quenched incubation mixtures was assayed by means of a radioimmunoassay, using components of a renin radioimmunoassay kit supplied by Becton Dickinson and Co. (Orangeburg, N.Y.). This radioimmunoassay was based upon the one developed by Haber et al., J. Clin. Endocrinol., 29, pp. 1349-1355 (1969).

The compounds of the present invention can be administered as antihypertensive agents by either the oral or parenteral routes of administration, with the former being preferred for reasons of patient convenience and comfort. In general, these antihypertensive compounds are normally administered orally in dosage ranging from about 0.1 mg to about 20 mg per kg of body weight per day and 0.1 mg to about 5 mg per kg of body weight per day when given parenterally; variations will necessarily occur depending upon the condition of the subject being treated and the particular compound being administered. Typically, treatment is commenced at a low daily dosage and increased by the physician only if necessary. It is to be noted that these compounds may be administered in combination with pharmaceutically acceptable carriers by either of the routes previously indicated, and that such administration can be carried out in both single and multiple dosages.

The novel compounds of the invention can be orally administered in a wide variety of different dosage forms, i.e., they may be formulated with various pharmaceutically acceptable inert carriers in the form of tablets, capsules, lozenges, troches, hard candies, powders, sprays, aqueous suspensions, elixirs, syrups and the like. Such carriers included solid diluents or fillers, sterile aqueous media and various non-toxic organic solvents, etc. Moreover, such oral pharmaceutical formulations can be suitably sweetened and/or flavored by means of various agents of the type commonly employed for such purposes. In general, the compounds of this invention are present in such oral dosage forms at concentration levels ranging from about 0.5% to about 90% by weight of the total composition, in amounts which are sufficient to provide the desired unit dosages.

For purposes of oral administration, tablets containing various excipients such as sodium citrate, calcium carbonate and calcium phosphate may be employed along with various disintegrants such as starch and preferably potato or tapioca starch, alginic acid and certain complex silicates, together with binding agents such as polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules; preferred materials in this connection would also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the essential active ingredient therein may be combined with various sweetening or flavoring agents, coloring matter or dyes and, if so desired, emulsifying and/or suspending agents as well, together with such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

The compounds of the present invention are also useful in the diagnosis of hypertension and congestive heart failure.

The following examples illustrate the invention but are not to be construed as limiting the same.

EXAMPLE 1

5 2R,4S,5S-6-Cyclohexyl-5-(5'-chloroindol-2'-ylcarbonylamino)-4-hydroxy-2-(2'-chloro-2'-propenyl)-N-methylhexanamide (HET = 5-chloroindol-2-yl; R₁ = cyclohexyl; R₂ = -C(Cl)=CH₂; and R₃ = CH₃)

10 1A. 2R,4S,5S-6-cyclohexyl-5-(5'-chloroindol-2'-ylcarbonylamino)-2-(2'-chloro-2'-propenyl)-gamma-hexanolactone

To 25 ml of methylene chloride at 0° C. was added 165.5 mg (0.5 mmol) of 2R,4S,5S-6-cyclohexyl-5-amino-2-(2'-chloro-2'-propenyl)-gamma-hexanolactone hydrochloride, 50.6 mg (0.5 mmol) of N-methyl morpholine, 97.8 mg (0.5 mmol) of 5-chloroindole-2-carboxylic acid, 67.5 mg (0.5 mmol) of N-hydroxybenzotriazole and 103 mg (0.5 mmol) of dicyclohexylcarbodiimide and the resulting reaction mixture stirred at room temperature overnight. The reaction mixture was filtered and the filtrate concentrated to dryness. The residue was redissolved in ethyl acetate, washed successively with water, a saturated sodium bicarbonate solution and brine and dried over magnesium sulfate. The solvent was removed and the residue, 280 mg, was chromatographed on silica gel using chloroform as the eluent to give 226 mg of the desired product.

25 1B. 2R,4S,5S-6-cyclohexyl-5-(5'-chloroindol-2'-ylcarbonylamino)-4-hydroxy-2-(2'-chloro-2'-propenyl)-N-methylhexanamide

A solution of 226 mg of the product of Example 1A in 10 ml of methanol was saturated with methyl amine gas the reaction mixture allowed to stand at room temperature for 1.5 hours. The reaction was concentrated to dryness and the residue triturated with ether to afford 155 mg of the desired product.

30 The NMR (CD₃OD) showed absorption at 2.7 (3H, s), 5.15 (2H, m) and 7.0-8.0 (4H, m) ppm.

EXAMPLE 2


35 Employing the procedure of Example 1A-1B, and starting with the appropriate starting reagents, the following compounds were prepared:

40

45



50



55



X	Y	R ₁	R ₂	R ₃	NMR (60 MHz) delta(CD ₃ OD)
5-Cl	H	C ₆ H ₁₁	i-C ₃ H ₇	CH ₃	0.95 (9H, d, J=5 Hz), 2.7 (3H, s), 7.0-8.0 (4H, m)
5-Cl	H	C ₆ H ₁₁	$\begin{array}{c} \text{-C=CH}_2 \\ \\ \text{CH}_3 \end{array}$	CH ₃	1.65 (3H, s), 2.7 3H, s), 4.65 (2H, m), 7.0-8.0 (4H, m)
5-Cl	H	C ₆ H ₁₁	-CH=CH ₂	CH ₃	2.7 (3H, s), 4.95 (2H, m), 7.0-8.0 (4H, m)
5-Cl	H	C ₆ H ₁₁	$\begin{array}{c} \text{-C=CH}_2 \\ \\ \text{Cl} \end{array}$	CH ₃	2.7 (3H, s), 5.15 (2H, m), 7.0-8.0 (4H, m)
5-Cl	H	C ₆ H ₁₁	$\begin{array}{c} \text{-C=CH}_2 \\ \\ \text{Cl} \end{array}$	H	5.15 (2H, m), 7.0- 8.0 (4H, m)
5-Cl	H	i-C ₃ H ₇	i-C ₃ H ₇	CH ₃	0.95 (12H, m), 2.7 (3H, s), 7.0-8.0 (4H, m)

	X	Y	R ₁	R ₂	R ₃	MMR (60 MHz) delta(CD ₃ OD)
5	H	H	C ₆ H ₁₁	$\begin{array}{c} -C=CH_2 \\ \\ Br \end{array}$	CH ₃	2.7 (3H, s), 5.5 (2H, m), 7.0-8.0 (5H, m)
10	5-Cl	H	C ₆ H ₁₁	$\begin{array}{c} -CH=CH \\ \\ CH_2N_3 \end{array}$	CH ₃	2.7 (3H, s), 7.0-8.0 (4H, m)
15	5-Cl	H	C ₆ H ₁₁	$\begin{array}{c} -CH-CH_3 \\ \\ OH \end{array}$	CH ₃	1.1 (3H, d, J=5 Hz), 2.7 (3H, s), 7.0-8.0 (4H, m)
20	5-Cl	H	C ₆ H ₁₁	$\begin{array}{c} -CH=CH \\ \\ Cl \end{array}$	CH ₃	2.7 (2H, m), 5.6-6.7 (2H, m), 7.0-8.0 (4H, m)
25	5-Cl	H	C ₆ H ₁₁	CH=C(CH ₃) ₂	CH ₃	1.8 (6H, br s), 2.7 (3H, s), 7.0-8.0 (4H, m)
30	3-Cl	H	C ₆ H ₁₁	i-C ₃ H ₇	CH ₃	0.95 (6H, d, J=5 Hz), 2.7 (3H, s)
35	5-Cl	H	C ₆ H ₁₁	$\begin{array}{c} -C=CH_2 \\ \\ Br \end{array}$	CH ₃	2.7 (3H, s), 5.5 (2H, m), 7.0-8.0 (4H, m)
40	5-Cl	H	C ₆ H ₁₁	i-C ₃ H ₇	CH ₃	0.95 (6H, d, J=5 Hz), 2.4 (3H, s), 2.7 (3H, s), 7.0-8.0 (4H, m)
45	5-Cl	H	C ₆ H ₁₁	$\begin{array}{c} CH_2CHC_2H_5 \\ \\ CH_3 \end{array}$	CH ₃	0.95 (12H, m), 2.7 (3H, s), 7.0-8.0 (4H, m)
50						
55						

	<u>X</u>	<u>Y</u>	<u>R₁</u>	<u>R₂</u>	<u>R₃</u>	NMR (60 MHz) <u>delta(CD₃OD)</u>
5	5-Cl	H	C ₆ H ₁₁	-CH=CH ₂	H	2.7 (3H, s), 4.95 (2H, m), 7.0-8.0 (4H, m)
10	5-Cl	H	C ₆ H ₁₁	-C ₆ H ₅	CH ₃	2.7 (3H, s), 7.0- 8.0 (9H, m)
15	H	H	C ₆ H ₁₁	i-C ₃ H ₇	CH ₃	0.95 (6H, d, J=5 Hz), 2.7 (3H, s), 7.0-8.0 (5H, m)
20	5-F	H	C ₆ H ₁₁	i-C ₃ H ₇	CH ₃	0.95 (6H, d, J=5 Hz), 2.7 (3H, s), 7.0-8.0 (4H, m)
25	H	H	C ₆ H ₁₁	-CH=CH ₂	CH ₃	2.7 (3H, s), 4.95 (2H, m), 7.0-8.0 (5H, m)
30	5-Cl	H	C ₆ H ₁₁	-CH=CHCl	-(CH ₂) ₂ 	2.6 (6H, m), 5.15 (2H, m), 6.9-7.7 (4H, m)
35	5-Cl	H	C ₆ H ₁₁	-CH=CHCH ₃	CH ₃	1.6 (3H, m), 2.7 (3H, s), 6.9-7.7 (4H, m)
40	5-Cl	H	C ₆ H ₁₁	-CH=CHCH ₃	-(CH ₂) ₂ 	1.6 (3H, m), 2.6 (6H, m), 5.3-5.7 (2H, m)
45						
50						
55						

	<u>X</u>	<u>Y</u>	<u>R₁</u>	<u>R₂</u>	<u>R₃</u>	NMR (60 MHz) delta(CD ₃ OD)
5	5-Cl	H	C ₆ H ₁₁	i-C ₃ H ₇	-(CH ₂) ₂ 	0.95 (6H, d, J=5 Hz), 2.7 (3H, s), 6.9-7.7 (4H, m)
10	5-Cl	H	C ₆ H ₁₁	-C(Cl)=CH ₂	-(CH ₂) ₂ 	2.6 (6H, m), 5.15 (2H, m), 5.9-7.7 (4H, m)
15						

EXAMPLE 3

2R,4S,5S-6-Cyclohexyl-5-(5'-chloroindol-3'-ylcarbonylamino)-4-hydroxy-2-(2'-methylpropyl)-N-methylhexanamide
(HET = 5-chloroindol-3-yl; R₁ = cyclohexyl; R₂ = -i-propyl; and R₃ = CH₃)

3A. 2R,4S,5S-6-cyclohexyl-5-(5'-chloroindol-3'-ylcarbonylamino)-2-(2'-methylpropyl)-gamma-hexanolactone

To 10 ml of methylene chloride at 0° C. was added 20 mg (0.1 mmol) of 5-chloroindole-3-carboxylic acid, 30 mg (0.1 mmol) of 2R,4S,5S-6-cyclohexyl-5-amino-2-(2'-methylpropyl)-gamma-hexanolactone hydrochloride, 15 μl (0.1 mmol) of diethyl cyanophosphate and 42 μl (0.3 mmol) of triethylamine. Dimethylformamide (1 ml) was added and the reaction mixture allowed to stir at room temperature overnight. The reaction was concentrated to dryness and redissolved in ethyl acetate which was washed successively with water (2x), a saturated sodium bicarbonate solution (2x) and a brine solution. The solution was dried over sodium sulfate and concentrated to an oil, 51 mg.


3B. 2R,4S,5S-6-cyclohexyl-5-(5' chloroindol-3'-ylcarbonylamino)-4-hydroxy-2-(2'-methylpropyl)-N-methylhexanamide

A solution of the product of Example 3A in 10 ml of methanol was saturated with methyl amine at room temperature and the resulting reaction mixture was stirred at room temperature for 2 hours. The reaction was evaporated to dryness giving 61 mg of a foam which was chromatographed on silica gel using methanol-chloroform as the eluent. The fractions containing the product were combined and concentrated to give 14 mg of the desired product.

The NMR (CD₃OD) showed absorption at 0.95 (6H, d, J = 5 Hz), 2.7 (3H, s) and 7.0-8.0 (4H, m) ppm.

EXAMPLE 4

Using the procedure of Example 3A-3B, and starting with the requisite starting reagents, the following compounds were prepared:



15

X	Y	R ₁	R ₂	R ₃	R ₄	NMR (60 MHz) delta (CD ₃ OD)
H	H	C ₆ H ₁₁	i-C ₃ H ₇	CH ₃	H	0.95 (6H, d, J=5 Hz), 2.7 (3H, s), 7.0-8.0 (5H, m)

20

7-CH ₃	H	C ₆ H ₁₁	i-C ₃ H ₇	CH ₃	H	0.95 (6H, d, J=5 Hz), 2.7 (3H, s), 2.4 (3H, s), 6.8-8.0 (4H, m)
-------------------	---	--------------------------------	---------------------------------	-----------------	---	---

25

30

5-Cl	H	C ₆ H ₁₁	i-C ₃ H ₇	CH ₃	H	0.95 (6H, d, J=5 Hz), 2.7 (3H, s), 7.0-8.0 (4H, m)
------	---	--------------------------------	---------------------------------	-----------------	---	--

35

X	Y	R ₁	R ₂	R ₃	R ₄	NMR (60 MHz) delta (CD ₃ OD)
5-Cl	H	C ₆ H ₁₁	-C-CH ₂ Cl	CH ₃	H	2.7 (3H, s) 5.15 (2H, m), 7.0-8.0 (4H, m)

40

45

H	H	C ₆ H ₁₁	i-C ₃ H ₇	CH ₃	CH ₃	0.95 (6H, d, J=5 Hz), 2.7 (3H, s), 3.7 (3H, s), 7.0-8.0 (4H, m)
---	---	--------------------------------	---------------------------------	-----------------	-----------------	---

50

55

5-CN	H	C ₆ H ₁₁	i-C ₃ H ₇	CH ₃	H	0.95 (6H, d, J=5 Hz), 2.7 (3H, s), 7.0-8.0 (4H, m)
------	---	--------------------------------	---------------------------------	-----------------	---	--

EXAMPLE 5

2R,4S,5S-6-Cyclohexyl-5-(5'-bromoindol-3'-ylacetyl-amino)-4-hydroxy-2-(2'-methylpropyl)-N-methylhexanamide
 (HET = 5-bromoindol-3-ylmethyl; R₁ = cyclohexyl; R₂ = i-propyl; R₃ = CH₃)

10 5A. 2R,4S,5S-6-cyclohexyl-5-(5'-bromoindol-3'-ylacetyl-amino)-2-(2'-methylpropyl)-gamma-hexanolactone

To 10 ml of methylene chloride and 0.3 ml of dimethylformamide at 0° C. was added 30 mg (0.1 mmol) of 2R,4S,5S-6-cyclohexyl-5-amino-2-(2'-methylpropyl)-gamma-hexanolactone hydrochloride, 25 mg (0.1 mmol) of 5-bromoindole-3-acetic acid, 21 mg (0.1 mmol) of dicyclohexylcarbodiimide, 14 mg (0.1 mmol) of N-hydroxybenzotriazole and 11 μ l (0.1 mmol) of N-methylmorpholine and the reaction mixture allowed to stir overnight at room temperature. The reaction was filtered and the filtrate evaporated to dryness. The residue was redissolved in ethyl acetate which was washed successively with water, a saturated sodium bicarbonate solution and a brine solution, and dried over sodium sulfate. The solvent was removed in vacuo and the residue, 54 mg, flash chromatographed on silica gel using 3% acetone as the eluent, 28 mg.

25 5B. 2R,4S,5S-6-cyclohexyl-5-(5'-bromoindol-3'-ylacetyl-amino)-4-hydroxy-2-(2'-methylpropyl)-N-methylhexanamide

A methanol (10 ml) solution containing 28 mg of the product of Example 5A was saturated with methyl amine at room temperature and allowed to stir for 2 hours. The reaction was evaporated to dryness and the residue triturated with hexane to give 30 mg of the desired product.

The NMR (CD₃OD) showed absorption at 0.95 (6H, d, J = 5 Hz), 2.7 (3H, s) and 7.0-8.0 (4H, m) ppm.

EXAMPLE 6


35 Employing the procedure of Example 5A-5B, and using the appropriate starting materials, the following compounds were prepared:

40

45

50

55



15

<u>X</u>	<u>Y</u>	<u>n</u>	<u>R₁</u>	<u>R₂</u>	<u>R₃</u>	NMR (60 MHz) delta (CD ₃ OD)
5-CH ₃	H	1	C ₆ H ₁₁	i-C ₃ H ₇	CH ₃	0.95 (6H, d, J=5 Hz), 2.7 (3H, s), 2.4 (3H, s), 7.0-8.0 (4H, m)

20

5-Br	H	1	C ₆ H ₁₁	i-C ₃ H ₇	CH ₃	0.95 (6H, d, J=5 Hz), 2.7 (3H, s), 7.0-8.0 (4H, s)
------	---	---	--------------------------------	---------------------------------	-----------------	--

25

<u>X</u>	<u>Y</u>	<u>n</u>	<u>R₁</u>	<u>R₂</u>	<u>R₃</u>	NMR (60 MHz) delta (CD ₃ OD)
H	H	2	C ₆ H ₁₁	i-C ₃ H ₇	CH ₃	0.95 (6H, d, J=5 Hz), 2.7 (3H, s)

30

35

H	H	1	C ₆ H ₁₁	i-C ₃ H ₇	CH ₃	0.95 (6H, d, J=5 Hz), 2.7 (3H, s)
---	---	---	--------------------------------	---------------------------------	-----------------	--------------------------------------

40

EXAMPLE 7

45 2R,4S,5S-6-Cyclohexyl-5-(6'-azaindol-2'-ylcarbonylamino)-4-hydroxy-2-(2'-methylpropyl)-N-methylhexanamide
 (HET = 6-azaindol-2-yl; R₁ = cyclohexyl; R₂ = i-propyl; R₃ = CH₃)

50 7A. 2R,4S,5S-6-cyclohexyl-5-(6'-azaindol-2'-ylcarbonylamino)-2-(2'-methylpropyl)-gamma-hexanolactone

To 10 ml of methylene chloride and 0.5 ml of dimethylformamide at 0° C. was added 37 mg (0.1 mmol) of 2R,4S,5S-6-cyclohexyl-5-amino-2-(2'-methylpropyl)-gamma-hexanolactone hydrochloride, 33 mg (0.2 mmol) of 6-azaindole-2-carboxylic acid, 25 mg (0.1 mmol) of dicyclohexylcarbodiimide, 17 mg (0.1 mmol) of N-hydroxybenzotriazole and 20 μl (0.1 mmol) of N-methylmorpholine, and the reaction allowed to stir at room temperature for 48 hours. The reaction was filtered, washed with water and a saturated sodium bicarbonate solution and dried over sodium sulfate. Evaporation of the solvent gave 55 mg of the desired product.


7B. 2R,4S,5S-6-cyclohexyl-5-(6'-azaindol-2'-ylcarbonylamino)-4-hydroxy-2-(2'-methylpropyl)-N-methylhexanamide

A solution of 55 mg of the product of Example 7A in 10 ml of methanol was saturated with methyl amine at room temperature and the reaction allowed to stand for 2 hours. The solvent was removed in vacuo and the residue chromatographed on silica gel using methanolchloroform as the eluent to give 9.7 mg of product.

The NMR spectrum (CD₃OD) showed absorption at 0.95 (6H, d, J=5 Hz) and 2.7 (3H, s) ppm.

EXAMPLE 8

Employing the procedure of Example 7A-7B, and starting with the appropriate starting materials, the following compounds were prepared:



	HET	NMR (60 MHz) delta (CD ₃ OD)
5		0.95 (6H, d, J=5 Hz), 2.7 (3H, s)
10		0.95 (6H, d, J=5 Hz), 2.7 (3H, s)
15		0.95 (6H, d, J=5 Hz), 2.7 (3H, s)
20		0.95 (6H, d, J=5 Hz), 2.7 (3H, s)

EXAMPLE 9

30 2R,4S,5S-6-Cyclohexyl-5-(indol-2'-ylcarbonylamino)-4-hydroxy-2-(4'-aminobutyl)-N-methylhexanamide
 (HET = indol-2-yl; R₁ = cyclohexyl; R₂ = (CH₂)₄NH₂; R₃ = CH₃)

35 A mixture of 75 mg of 2R,4S,5S-6-cyclohexyl-5-(5'-chloroindol-2'-ylcarbonylamino)-4-hydroxy-2-(4'azido-2'-butenyl)-N-methylhexanamide, 35 mg of palladium-on-charcoal and 5 ml of acetic acid in 5 ml of methanol was shaken in a hydrogen atmosphere at 50 psi for 6 hours. An additional 35 mg of catalyst was added and the reduction continued overnight. The catalyst was filtered and the filtrate concentrated in vacuo to give 66 mg of product. The residue was triturated with ether and filtered, 50 mg.

40 The NMR (CD₃OD) spectrum showed absorption at 2.7 (3H, s) and 7.0-8.0 (5H, m) ppm.

PREPARATION A

45 2R,4S,5S-6-Cyclohexyl-5-(t-butoxycarbonylamino)-2-(2'-chloro-2'-propenyl)-gamma-hexanolactone








50 To a tetrahydrofuran solution containing 37.5 mmol of lithium diethylamide at -70° C. (prepared from 23.4 ml of 1.6M butyl lithium hexane and 4.26 g of diethyl amine in 50 ml of dry tetrahydrofuran) was added dropwise a solution of 4.67 g (15 mmol) of 4S,5S-6-cyclohexyl-5-(5-butoxycarbonylamino)-gamma hexanolactone in 25 ml of tetrahydrofuran. After 30 minutes at -78° C. a solution of 3.64 g (16 mmol) of 2-chloro-3-iodopropene in 25 ml of tetrahydrofuran was added dropwise at -70° C. After 2 hours the reaction mixture was quenched with 10 ml of a saturated ammonium chloride solution added dropwise at -78° C., and the resulting mixture allowed to warm to room temperature. The solvent was removed in vacuo and the residue extracted with diethyl ether. The ether solution was washed with a 10% citric acid solution, a saturated sodium bicarbonate solution and a brine solution. The ether solution was then dried over magnesium sulfate and concentrated to give 6.83 g of an oil, which was chromatographed on silica gel

using ethyl acetate-hexane as the eluent. The fractions containing the product were combined and concentrated to give 2.38 g of the desired product.

The NMR spectrum (CDCl_3) showed absorption at 1.4 (9H, s) ppm.

PREPARATION B

Employing the procedure of Preparation A, and using the appropriate starting reagents, the following intermediates were synthesized:



PREPARATION C

2R,4S,5S-6-Cyclohexyl-5-(t-butoxycarbonylamino)-2-(4'-azido-2-butenyl)-gamma-hexanolactone

A solution of 710 mg (1.6 mmol) of 2R,4S,5S-6-cyclohexyl-5-(t-butoxycarbonylamino)-2-(4'-bromo-2-butenyl)-gamma-hexanolactone and 986 mg (15.2 mmol) of sodium azide in 75 ml of dimethylsulfoxide-water (2:1; v:v) was allowed to stir overnight at room temperature. The reaction was poured into 500 ml of water and the product extracted with ethyl acetate. The extracts were combined and washed successively with water and a brine solution, and were dried over magnesium sulfate. The solvent was removed in vacuo to give 73 mg of the desired product.

PREPARATION D5 2R,4S,5S-6-Cyclohexyl-5-(t-butoxycarbonylamino)-2-(2'-hydroxypropyl)-gamma-hexanolactone

A solution of 100 mg (0.27 mmol) of 2R,4S,5S-6-cyclohexyl-5-(t-butoxycarbonylamino)-2-(2'-methyl-2'-propenyl)-gamma-hexanolactone in 15 ml of methylene chloride was saturated with ozone at -78° C. until
10 there was a permanent blue coloration. The reaction mixture was allowed to stand for 30 minutes at -78° C. and was then purged of excess ozone at -78° C. with a rapid stream of nitrogen. Tetrabutyl ammonium borohydride (140 mg, .54 mmol) was added at -78° C. and the reaction mixture was allowed to stand at 0° C. for 2 days. An additional 140 mg of borohydride was added and the reaction mixture allowed to stand
15 overnight at room temperature. The solution was washed with water and a brine solution, dried over sodium sulfate and evaporated to give 250 mg of crude product. Flash chromatographing using chloroform-methanol (99:1, v:v) afforded 45 mg of the desired intermediate.

The NMR (CDCl₃) spectrum showed absorption at 1.4 (9H, s) ppm.

20

PREPARATION E25 2R,4S,5S-6-Cyclohexyl-5-amino-2-(2'-chloro-2'-propenyl)-gamma-hexanolactone hydrochloride

25

A solution of 385 mg (1 mmol) of 2R,4S,5S-6-cyclohexyl-5-(t-butoxycarbonylamino)-2-(2'-chloro-2'-propenyl)-gamma-hexanolactone in 10 ml of 4.7N hydrogen chloride in dioxane was allowed to stir for 2
30 hours at room temperature. The solvent was removed in vacuo to give 331 mg of the desired amine hydrochloride.

PREPARATION F

35









Using the procedure of Preparation E, and employing the intermediates of Preparations A, B and D, the following intermediates were prepared:

40

45


50

55




Claims


1. A compound selected from those of the formula



55 or a pharmaceutically acceptable salt thereof, wherein HET is




or a group of the formula



where X is hydrogen, (C₁-C₃)alkyl, (C₁-C₃)alkoxy, fluoro, chloro, bromo or cyano; Y is hydrogen, (C₁-C₃)alkyl, (C₁-C₃)alkoxy, fluoro or chloro; R₄ is hydrogen or (C₁-C₃)alkyl; n is an integer of 0 to 2; R₁ is (C₆-C₈)cycloalkyl or i-propyl; R₂ is (C₃-C₅)alkyl, phenyl, methylvinyl, dimethylvinyl, halovinyl, hydroxy(C₁-C₃)alkyl or amino(C₁-C₄)alkyl; and R₃ is (C₁-C₆) or morpholinoethyl.

2. A compound of claim 1, wherein HET is a group of the formula




R₁ is cyclohexyl and R₃ is methyl.

3. The compound of claim 2, wherein X is 5-chloro, Y is hydrogen, R₄ is hydrogen, n is 0 and R₂ is i-propyl.

4. The compound of claim 2, wherein X and Y are each hydrogen, R₄ is methyl, n is 0 and R₂ is i-propyl.

5. A compound of claim 1, wherein HET is a group of the formula




R₁ is cyclohexyl and R₃ is methyl.

6. The compound of claim 5, wherein X is 5-chloro, Y is hydrogen, R₄ is hydrogen, n is 0 and R₂ is -CH(OH)CH₃.

7. The compound of claim 2, wherein X is 5-chloro, Y is hydrogen, R₄ is hydrogen, n is 0 and R₂ is -C-(Cl)=CH₂.

8. A compound of claim 1, wherein HET is a group of the formula



R₁ is cyclohexyl and R₂ is methylvinyl.

9. The compound of claim 8, wherein X is 5-chloro, Y is hydrogen, R₄ is hydrogen, n is 0, R₂ is -CH=CHCH₃ and R₃ is methyl.


10. The compound of claim 8, wherein X is 5-chloro, Y is hydrogen, R₄ is hydrogen, n is 0, R₂ is -CH=CHCH₃ and R₃ is morpholinoethyl.

11. A pharmaceutical composition comprising an antihypertensive effective amount of a compound according to claim 1 and a pharmaceutically acceptable diluent or carrier.


12. A compound as claimed in claim 1 for use in treating hypertension.

Claims for the following Contracting States: ES, GR


1. A process for preparing a compound of the formula




or a pharmaceutically acceptable salt thereof, wherein HET is



or a group of the formula



where X is hydrogen, (C_1-C_3) alkyl, (C_1-C_3) alkoxy, fluoro, chloro, bromo or cyano; Y is hydrogen, (C_1-C_3) alkyl, (C_1-C_3) alkoxy, fluoro or chloro; R_4 is hydrogen or (C_1-C_3) alkyl; n is an integer of 0 to 2; R_1 is (C_6-C_8) cycloalkyl or i -propyl; R_2 is (C_3-C_5) alkyl, phenyl, methylvinyl, dimethylvinyl, halovinyl, hydroxy (C_1-C_3) alkyl or amino (C_1-C_4) alkyl; and R_3 is (C_1-C_6) alkyl or morpholinoethyl characterized by reacting a compound of the formula



with an amine of the formula R_3NH_2 in a lower alcohol at room temperature until substantially complete and, optionally, reducing the compound wherein R_2 is omega- $N_3(C_2-C_4)$ alkylene or N_3CH_2- with hydrogen and palladium-on-charcoal to give the compound wherein R_2 is amino (C_1-C_4) alkyl.

(9)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 321 192 A3

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 88311798.8

(22) Date of filing: 14.12.88

(51) Int. Cl.⁵: **C07D 209/42**, C07D 471/04,
C07D 215/56, C07D 241/44,
C07D 207/34, C07D 209/18,
C07D 209/08, A61K 31/40,
A61K 31/47, A61K 31/495,
A61K 31/44, //(C07D471/04,
221:00,209:00),(C07D471/04,
235:00,221:00)

(30) Priority: 15.12.87 US 132373

(43) Date of publication of application:
21.06.89 Bulletin 89/25

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

(88) Date of deferred publication of the search report:
30.01.91 Bulletin 91/05


(71) Applicant: **PFIZER INC.**
235 East 42nd Street
New York, N.Y. 10017(US)

(72) Inventor: **Rosati, Robert Louis**
RD 3, Box 66A Deans Mill Road
Stonington Connecticut(US)

(74) Representative: **Moore, James William, Dr.**
Pfizer Limited Ramsgate Road
Sandwich Kent CT13 9NJ(GB)

(54) **Non-peptidic renin inhibitors.**

(57) Simple amides and derivatives thereof useful for inhibiting the angiotensinogen-cleaving action of the enzyme renin of the formula:



wherein R₁ is (C₆-C₈)cycloalkyl or i-propyl; R₂ is (C₃-C₅)alkyl, phenyl, methylvinyl, dimethylvinyl, halovinyl, hydroxy(C₁-C₃)alkyl or amino(C₁-C₄)alkyl; R₃ is (C₁-C₆)alkyl or morpholinoethyl; and Het is defined to include certain mono and bicyclic nitrogen-containing heterocyclic rings or heterocyclalkyl groups.

EP 0 321 192 A3



European
Patent Office

EUROPEAN SEARCH REPORT

Application Number

EP 88 31 1798

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
A	EP-A-0 184 550 (CIBA-GEIGY) " Examples 4,7,14,16,51,53 "	1,12	C 07 D 209/42 C 07 D 471:04 C 07 D 215:56
A	EP-A-0 181 110 (KISSEI PHARMACEUTICAL) " Abstract "	1,12	C 07 D 241:44 C 07 D 207:34 C 07 D 209/18 C 07 D 209/08 A 61 K 31:40 A 61 K 31:47 A 61 K 31:495 A 61 K 31:44 (C 07 D 471:04 C 07 D 221:00 C 07 D 209:00) (C 07 D 471:04 C 07 D 235:00 C 07 D 221:00)
			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
			C 07 D 209/00 C 07 D 471:00 C 07 D 215/00 C 07 D 241/00 C 07 D 207/00
The present search report has been drawn up for all claims			
Place of search		Date of completion of search	Examiner
The Hague		16 November 90	DE JONG B.S.
CATEGORY OF CITED DOCUMENTS			
X: particularly relevant if taken alone			
Y: particularly relevant if combined with another document of the same category			
A: technological background			
O: non-written disclosure			
P: intermediate document			
T: theory or principle underlying the invention			
E: earlier patent document, but published on, or after the filing date			
D: document cited in the application			
L: document cited for other reasons			
&: member of the same patent family, corresponding document			

